

Criterion for local distinguishability of arbitrary bipartite orthogonal states

Ping-Xing Chen^{1,2*} and Cheng-Zu Li¹

*1. Department of Applied Physics, National University of
Defense Technology,
Changsha, 410073,
P. R. China.*

*2. Laboratory of Quantum Communication and Quantum Computation,
University of Science and Technology of
China, Hefei, 230026, P. R. China
(February 1, 2008)*

Abstract

In this paper we present a necessary and sufficient condition of distinguishability of bipartite quantum states. It is shown that the operators to reliably distinguish states need only rounds of projective measurements and classical communication. We also present a necessary condition of distinguishability of bipartite quantum states which is simple and general. With this condition one can get many cases of indistinguishability. The conclusions may be useful in understanding the essence of nonlocality and calculating the distillable entanglement and the bound of distillable entanglement.

PACS number(s): 03.67.-a, 03.65.ud

Typeset using REVTeX

*E-mail: pxchen@nudt.edu.cn

One of the interesting features of non-locality in quantum mechanics is that a set of orthogonal quantum states cannot be distinguished if only a single copy of these states is provided and only local operations and classical communication (LOCC) are allowed, in general. The procedure of distinguishing quantum states locally is: Alice and Bob hold a part of a quantum system, which occupies one of m possible orthogonal states $|\Psi_1\rangle, |\Psi_2\rangle, \dots, |\Psi_i\rangle, \dots, |\Psi_m\rangle$. Alice and Bob know the precise form of these states, but don't know which of these possible states they actually hold. To distinguish these possible states they will perform some operations locally: Alice (or Bob) first measures her part. Then she tell the Bob her measurement result, according to which Bob measure his part. With the measurement results they can exclude some possibilities of the system [1]. Briefly speaking, the procedure of distinguishing quantum states locally is to exclude all or some possibilities by measurement on the system. Many authors have considered some schemes for distinguishing locally between a set of quantum states [1–7], both inseparable and separable. Bennett et al showed that some orthogonal product states cannot be distinguished by LOCC [2]. Walgate et al showed that any two states can be distinguished by LOCC [1]. For two-qubit systems (or $2 \otimes 2$ systems), any three of the four Bell states:

$$\begin{aligned} |A_1\rangle &= \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle) \\ |A_2\rangle &= \frac{1}{\sqrt{2}}(|00\rangle - |11\rangle) \\ |A_3\rangle &= \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle) \\ |A_4\rangle &= \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle) \end{aligned} \tag{1}$$

cannot be distinguished by LOCC if only a single copy is provided [4]. The distinguishability of quantum states has some close connections [8] with distillable entanglement [9] and the information transformation [10]. On one hand, using the upper bound of distillable entanglement, relative entropy entanglement [11] and logarithmic negativity [12], the authors in Ref [4] proved that some states are indistinguishable. On the other hand, using the rules on distinguishability one can discuss the distillable entanglement [8]. So the further analysis for distinguishability is meaningful. In this paper, we will give a necessary and sufficient condition of distinguishability of bipartite quantum states. It is shown that the operators to reliably distinguish states need only rounds of projective measurements and classical communication. We also present a necessary condition of distinguishability of bipartite quantum states which is simple and general. With this condition one can get many cases of indistinguishability.

Consider m possible orthogonal states shared between Alice and Bob. Any protocol to distinguish the m possible orthogonal states can be conceived as successive rounds of

measurement and communication by Alice and Bob. Let us suppose Alice is the first person to perform a measurement (Alice goes first [3]), and the first round measurement by Alice can be represented by operators $\{A_{1_j}\}$, where $A_{1_j}^+ A_{1_j}$ is known as a POVM element realized by Alice [13,14], and $\sum_j A_{1_j}^+ A_{1_j} = I$. If the outcome 1_j occurs, then the given $|\Psi\rangle$ becomes $A_{1_j} |\Psi\rangle$, up to normalization. After communicating the result of Alice's measurement to Bob, he carries out a measurement and obtain outcome 1_k . The given possible state $|\Psi\rangle$ becomes $A_{1_j} \otimes B_{1_k}(1_j) |\Psi\rangle$, where $B_{1_k}(1_j)$ is an arbitrary measurement operator of Bob which depend on the outcome 1_j of Alice's measurement. After N rounds of measurements and communication, there are many possible outcomes which corresponding to many measurement operators acting on the Alice and Bob's Hilbert space. Each of these operators is a product of the N sequential and relative operators $A_{1_j} \otimes B_{1_k}(1_j) A_{2_j}(1_j, 1_k) \otimes B_{2_k}(1_j, 1_k, 2_j) \dots A_{N_j}(1_j, 1_k, \dots, (N-1)_j) \otimes B_{N_k}(1_j, 1_k, \dots, (N-1)_j, N_k)$ carried out by Alice and Bob. We denote these operators as $\{A_i \otimes B_i\}$, where, $A_i \otimes B_i$ denotes one of these operators, which represent the effects of the N measurements and communication. If the outcome i occurs, the given $|\Psi\rangle$ becomes:

$$A_i \otimes B_i |\Psi\rangle \quad (2)$$

The probability p_i Alice and Bob gain outcome i is

$$p_i = \langle \Psi | A_i^+ \otimes B_i^+ A_i \otimes B_i | \Psi \rangle, \quad (3)$$

and

$$\sum_i A_i^+ \otimes B_i^+ A_i \otimes B_i = I \quad (4)$$

Suppose we define:

$$E_i = A_i^+ \otimes B_i^+ A_i \otimes B_i, \quad (5)$$

then E_i is a positive operator and that $\sum_i E_i = I$. E_i is similar to the POVM element $A_i^+ A_i$. We can regard E_i as a generalized POVM (GPOVM) element, which has same property as known POVM element. In fact, A_i can be written in the form $A_i = U_{A2} f_{A_i} U_{A1}$ [14], where f_{A_i} is a diagonal positive operator, U_{A2}, U_{A1} are two unitary operators, and similarly for B_i . If each of N Alice's operators denoted by A_i and each of N Bob's operators denoted by B_i are projectors, the final operators $A_i \otimes B_i$ are also projectors, i.e., $A_i \otimes B_i A_j \otimes B_j = \delta_{ij} A_i \otimes B_i$, and $\{E_i\}$ is a set of projective measurement.

The discuss above means that: whatever Alice and Bob choose to do, including they decide to involve an ancillary system; they perform local unitary operators and measurement; they use one-way or two-way communication, and do many rounds of measurements and communication, their final actions will be described by a set positive operators $\{E_i\}$. The probability of a given possible state $|\Psi\rangle$ yielding a certain outcome i is $\langle \Psi | A_i^+ \otimes B_i^+ A_i \otimes B_i | \Psi \rangle$.

Since a GPOVM element E_i has similar property as a POVM element, E_i can be represented in the form

$$E_i = (a_1^i |\varphi_1^i\rangle_A \langle\varphi_1^i| + \cdots + a_{m_a}^i |\varphi_{m_a}^i\rangle_A \langle\varphi_{m_a}^i| + \cdots) \otimes (b_1^i |\eta_1^i\rangle_B \langle\eta_1^i| + \cdots + b_{m_b}^i |\eta_{m_b}^i\rangle_B \langle\eta_{m_b}^i| + \cdots) \quad (6)$$

$$0 \leq a_{m_a}^i \leq 1, 0 \leq b_{m_b}^i \leq 1; 1 \leq m_a \leq N_a, 1 \leq m_b \leq N_b \quad (7)$$

where $\{|\varphi_{m_a}^i\rangle\}, \{|\eta_{m_b}^i\rangle\}$ is a set of bases of Alice's and Bob's, respectively; N_a, N_b is the dimensions of Alice's and Bob's Hilbert space, respectively.

Theorem 1. If a set of m orthogonal states $\{|\Psi_i\rangle\}$ is reliably locally distinguishable, if and only if there is a set of locally distinguishable product vectors, $\{|\varphi_i^{j_i}\rangle_A |\eta_i^{j_i}\rangle_B\}$, such that each state $|\Psi_i\rangle$ can be written as a superposition of some different product vectors in the set of these product vectors:

$$|\Psi_i\rangle = c_i^1 |\varphi_i^1\rangle_A |\eta_i^1\rangle_B + \cdots + c_i^{n_i} |\varphi_i^{n_i}\rangle_A |\eta_i^{n_i}\rangle_B \quad (8)$$

where $i = 1, \dots, m; j_i = 1, \dots, n_i$. n_i is the number of product bases in the state $|\Psi_i\rangle$

Proof: The proof of sufficiency is obviously. Now we prove the necessity. If a set of states is reliably locally distinguishable, there must be a set of GPOVM element $\{E_i\}$ such that if every outcome i occurs Alice and Bob know with certainty that they were given the state $|\Psi_i\rangle$. In a simple way we say that E_i can and only can "indicate" $|\Psi_i\rangle$.

Since each E_i only indicate a state $|\Psi_i\rangle$, the rank of E_i should be less than $N_a N_b$. Otherwise, E_i will indicates all states $\{|\Psi_i\rangle\}$. Without loss of generality, we suppose $a_1^i, \dots, a_{m_a}^i, b_1^i, \dots, b_{m_b}^i$ in Eq. (6) are nonzero, the other coefficient are zero, then the state $|\Psi_i\rangle$ should have all or part of the component

$$|\varphi_1^i\rangle_A |\eta_1^i\rangle_B, \dots, |\varphi_1^i\rangle_A |\eta_{m_b}^i\rangle_B, \dots, |\varphi_{m_a}^i\rangle_A |\eta_1^i\rangle_B, \dots, |\varphi_{m_a}^i\rangle_A |\eta_{m_b}^i\rangle_B. \quad (9)$$

The state $|\Psi_j\rangle$ ($i \neq j$) should have not these component. Otherwise, E_i will indicates more than one state. The effect of the operator E_i is project out the component

$$|\varphi_1^i\rangle_A |\eta_1^i\rangle_B, \dots, |\varphi_1^i\rangle_A |\eta_{m_b}^i\rangle_B, \dots, |\varphi_{m_a}^i\rangle_A |\eta_1^i\rangle_B, \dots, |\varphi_{m_a}^i\rangle_A |\eta_{m_b}^i\rangle_B \quad (10)$$

which only belongs to $|\Psi_i\rangle$. Because of the completeness of $\{E_i\}$ (which assures that each component in all possible states can be indicated by a GPOVM element) and the necessity of reliably distinguishing the possible states (which asks a GPOVM element only indicate the component only belonging to a possible states), the m possible states can be divided into many component each of which can be indicatd by a GPOVM element. It is to say that these component are locally distinguishable. If a operator E_i only indicate a state, then E_i can be replaced by a set of operators

$$E_{i1} = |\varphi_1^i\rangle_A \langle \varphi_1^i| \otimes |\eta_1^i\rangle_B \langle \eta_1^i|; \dots; \quad (11)$$

$$E_{im_b} = |\varphi_1^i\rangle_A \langle \varphi_1^i| \otimes |\eta_{m_b}^i\rangle_B \langle \eta_{m_b}^i|; \dots; \quad (12)$$

$$E_{im_a m_b} = |\varphi_{m_a}^i\rangle_A \langle \varphi_{m_a}^i| \otimes |\eta_{m_b}^i\rangle_B \langle \eta_{m_b}^i|,$$

each of which also only indicates the same state as E_i does. The effect of each operator E_{ij} ($j = 1, \dots, m_a m_b$) is to project out a component which only belong to a single state, for example, operator project out the component $|\varphi_1^i\rangle_A |\varphi_1^i\rangle_B$. Thus all product vectors which only belong to a state can be indicated by a set of operators $\{E_{ij}\}$, so all the product vectors are locally distinguishable and then are orthogonal. Because of the completeness of $\{E_i\}$ and the necessity of reliably distinguishing the possible states, each state $|\Psi_i\rangle$ must be the superposition of some locally distinguishable orthogonal product bases. This end the proof.

From theorem 1 it follows that if a set of operators $\{E_i\}$ can distinguish a set of states, there is a set of orthogonal projective operators $\{E_{ij}\}$ which can achieve the distinguishing the possible states. These operators can always be carried out by rounds of projective measurements and classical communication. It is to say that if a set of m states are distinguishable, the operator to distinguish the states can be always described as: First, Alice and Bob choose a person to go first to do measurement with her or his a set of orthogonal bases; After measurement, their Hilbert space collapse into a subspace which is orthogonal to the subspace discarded. According to the outcome, they know that the m possible states collapses into n' ($n' \leq n$) distinguishable new states, and choose a person to do measurement with her or his other set of orthogonal bases once more, and so on. After many rounds of projective measurements and classical communication, they may get a final state which only belongs to one of the possible states, and the Hilbert space collapses into a lower subspace. The final state may be entangled state or separable state. If the final state is entangled one can choose end of the operators, also can choose continue to do measurement such that it collapses into one of its orthogonal product states. In each round of measurement and communication, Alice and Bob must choose an appropriate person to do measurement. The different round of operator many need different person to do first, in general. For example, six states in a $4 \otimes 4$ system,

$$|\Psi_1\rangle = |0\rangle_A |0\rangle_B; |\Psi_2\rangle = |1\rangle_A (|0\rangle + |1\rangle)_B; |\Psi_3\rangle = |0\rangle_A |1\rangle_B + |1\rangle_A (|0\rangle - |1\rangle)_B \quad (13)$$

$$|\Psi_4\rangle = |2\rangle_A |2\rangle_B; |\Psi_5\rangle = (|2\rangle + |3\rangle)_A |3\rangle_B; |\Psi_6\rangle = |3\rangle_A |2\rangle_B + (|2\rangle - |3\rangle)_A |3\rangle_B \quad (14)$$

one can first choose to do measurement with Alice's bases

$$E_1 = |0\rangle_A \langle 0| + |1\rangle_A \langle 1|; \quad E_2 = |2\rangle_A \langle 2| + |3\rangle_A \langle 3|$$

if the outcome is E_1 they must choose Alice to do first to do the second measurement; if the outcome is E_2 they must choose Bob to go first to do the second measurement. The distinguishability of states in $2 \otimes n$ systems is a special example in which after Alice do measurement, the possible states collapse into some orthogonal product bases, so the possible states can be written as the form in the Theorem 1 of Walgate's paper [3].

Before giving theorem 2 in this paper, we define a concept of *Schmidt number*. If a pure state $|\Psi\rangle$ have following Schmidt decomposition:

$$|\Psi\rangle = \sum_{i=1}^l \sqrt{P_i} |\nu_i\rangle_A |\eta_i\rangle_B, \quad \sum_{i=1}^l P_i = 1 \quad (15)$$

where $|\nu_i\rangle_A$'s and $|\eta_i\rangle_B$'s are orthogonal bases of Alice and Bob, respectively, we say $|\Psi\rangle$ has *Schmidt number* l . Keep this in mind we will start from the following theorem.

Theorem 2: Alice and Bob share a quantum system, which possesses one of pure orthogonal states $|\Psi_1\rangle, |\Psi_2\rangle, \dots, |\Psi_i\rangle, \dots, |\Psi_m\rangle$. If the dimensions of Hilbert space of Alice's part and Bob's part are N_a and N_b , respectively, one cannot distinguish deterministically a set of orthogonal states for which the sum of *Schmidt numbers* is more than $N_a N_b$ when only a single copy is provided.

Theorem 2 can be expressed briefly as: one cannot distinguish a set of orthogonal states the sum of Schmidt number of which is more than the dimensions of whole Hilbert space of the quantum system.

From theorem 2 one can get the following interesting cases:

Case1: For $n \otimes n$ systems one cannot distinguish deterministically $n + 1$ states, each of which has Schmidt number n . For example, one can at most distinguish two entangled states in $2 \otimes 2$ systems.

Case 2: For $n \otimes n$ systems, if one can distinguish n^2 orthogonal states, these states must be orthogonal bases .

Proof of theorem 2: If the m possible orthogonal states are reliably locally distinguishable there are a set of orthogonal product basis each of which only emerges in one state of the m possible states, and therefore only "indicates" one state. A pure state with Schmidt numbers l includes at least l orthogonal product bases, so if the sum of Schmidt numbers of the m possible states is more than $N_a N_b$, these possible states surely include product bases more than $N_a N_b$. However, the number of orthogonal product bases in a $N_a \otimes N_b$ system is at most $N_a N_b$. So these m possible states with Schmidt number more than $N_a N_b$ cannot be written as the form in theorem 1, which means they are not distinguishable. This completes the proof theorem 2.

According to the theorem 2 we can discuss completely the case for $2 \otimes 2$ systems in which there are at most four orthogonal states, as be shown in Ref [3]. Four orthogonal states

can be distinguished deterministically if and only if they are four product states. This is obviously from the theorem above or case 2; three orthogonal states can be distinguished deterministically if and only if almost one state is entangled state and the other states are product states. This is because any two product orthogonal states can be written as: $|\nu_1\rangle|\eta_1\rangle$ and $|\nu_2\rangle|\eta_2\rangle$, where at least one of the inner product $\langle\nu_1|\nu_2\rangle$ and $\langle\eta_1|\eta_2\rangle$ is zero. Without loss of generality, we let $\langle\nu_1|\nu_2\rangle = 0$, then $|\nu_1\rangle, |\nu_2\rangle$ consist of a complete set of orthogonal bases in two dimensions Hilbert space. Any entangled state in $2 \otimes 2$ systems can be expressed as: $a|\nu_1\rangle|\eta'_1\rangle + b|\nu_2\rangle|\eta'_2\rangle$. As these three states are orthogonal, the following relation should hold: $\langle\eta_1|\eta'_1\rangle = \langle\eta_2|\eta'_2\rangle = 0$. This means the three states are distinguishable deterministically. If more than two states of the three orthogonal states are entangled, the sum of their Schmidt numbers would be more than four, which does not satisfy the necessary condition for distinguishability in the theorem above. So if and only if not less than two states of three orthogonal states are product states, the three states are distinguishable.

In summary, we present a necessary and sufficient condition of distinguishability of bipartite quantum states. It is shown that the operators to reliably distinguish states need only rounds of projective measurements and classical communication.. We also present a necessary condition of distinguishability of bipartite quantum states which is simple and general. With this condition one can get many cases of indistinguishability. These results come directly from the limits on local operations, not from the upper bound of distillable entanglement [4], So we believe that they may be useful in calculating the distillable entanglement or the bound of distillable entanglement. The further works may be the applications of these results.

ACKNOWLEDGMENTS

We would like to thank professor M.B. Plenio for his helpful suggestions and professor Guangcan Guo for his help to this work.

REFERENCES

- [1] J.Walgate, A.J.Short, L.Hardy and V.Vedral, Phys.Rev.Lett.85,4972 (2000).
- [2] C.H. Bennett, D.P. DiVincenzo, C.A. Fuchs, T.Mor, E.Rains, P.W. Shor, J.A. Smolin, and W.K. Wootters, Phys. Rev. A 59,1070 (1999) or quant-ph/9804053.
- [3] J. Walgate and L. Hardy, quant-ph/0202034, to published in Phys. Rev. Lett
- [4] S.Ghosh, G.Kar, A.Roy, A.Sen and U.Sen, Phys.Rev.Lett.87, 277902 (2001); S. Ghosh, G.Kar, A.Roy, D.Sarkar, A.Sen(De) and U.Sen, quant-ph/0111136 (2001)
- [5] S. Virmani, M.F. Sacchi, M.B. Plenio and D. Markham, Physics Letters A 288, 62-68 (2001);
- [6] M. Horodecki, P. Horodecki, and R. Horodecki, Acta Physica Slovaca, 48, (1998) 141, or quant-ph/9805072
- [7] Y.-X.Chen and D.Yang, Phys.Rev.A 64, 064303 (2001);
- [8] P.-X Chen and C.-Z Li, quant-ph/0202165
- [9] C. H. Bennett, G. Brassard, S. Popescu, B. Schumacher, J. A. Smolin, and W. K. Wootters, Phys. Rev. Lett **76** 722 (1996); C. H. Bennett, D. P. Divincenzo, J. A. Smolin, and W. K. Wootters, Phys. Rev. A **54**, 3824 (1996)
- [10] J. Eisert, T. Felbinger, P. Papadopoulos, M.B. Plenio and M. Wilkens, Phys. Rev. Lett. 84, 1611 (2000); L.Henderson and V.Vedral, Phys.Rev.Lett 84, 2263 (2000).
- [11] V.Vedral, M.B.Plenio, M.A.Rippin and P.L.Knight, Phys.Rev.Lett. 78, 2275 (1997); V.Vedral and M.B.Plenio, Phys.Rev.A 57, 1619 (1998)
- [12] G.Vidal and R.F.Werner, quant-ph/0102117 (2001);
- [13] H. Barnum, M. A. Nielsen and B. Schumacher, Phys.Rev.A 57, 4153 (1998);
- [14] N. Linden, S. Massar and S. Popescu, Phys.Rev.Lett 81, 3279 (1998)